

Introduction

On-line (recursive) estimation consists in sequential update of the statistics and if needed ongoing computation of the point estimates of the parameters. Each step, the newly measured data are added and used for improving the estimates. The process starts with a prior statistics to be recomputed at the first step. If there is no prior information the prior statistics can be constructed so they give uniform prior distribution of parameters (i.e. each value of them is equally probable). However, it is highly unreasonable, from the following reasons:

1. Always, there is some prior information - stronger or weaker, but it is.
2. With uniform prior density of parameters the initial estimates can be very wild and for example if the model is used for control, they can be the cause of the failure of the whole control process.
3. At the beginning of estimation we use only few data (at the first step only one data record). This record is under uncertainty and thus it can determine parameters that are far from the true values (imagine e.g. tossing a coin). Then the estimates must be dragged back (with partially fixed statistics and they do not need to reach the true positions).

To prevent these inconveniences, we can

- set the initial statistics so that they produce reasonable point estimates of the parameters,
- tighten the initial estimates appropriately so that they hold on for a while and can be moved only by a larger portion of data.

These two steps are universal (for arbitrary model), but for each one the way how to do it is a bit different. We shall demonstrate the way on the categorical model describing tossing of a coin.

Example

Let us have a discrete variable $y_t \in \{0, 1\}$, $t = 1, 2, \dots, N$ describing tossing a coin. The value $y_t = 1$ represents the result "head" and $y_t = 0$ is "tail". We have tossed the coin 5 times ($N=5$) and obtained the data

1 1 0 1 0

Perform step by step the estimation of the probability P ("head"). We know that in reality the probability is $P = 0.5$.


The estimation can be performed according to the statistical definition of probability

$$P = \frac{\text{number of heads}}{\text{number of experiments}}$$

which is listed in the following table (\hat{p} is the estimate of the probability)

t	numb. of heads	numb. of experiments	\hat{p}
1	1	1	1
2	2	2	1
3	2	3	$\frac{2}{3}$
4	3	4	$\frac{3}{4}$
5	3	5	$\frac{3}{5}$

We can see that at the beginning the estimates are totally wrong and only gradually they are refined. The estimates evolution is



Now, with our knowledge (that the coin is correct), we can construct so called fictitious prior data expressing our knowledge, e.g.


1 0

and use them before the estimation starts. We obtain

t	numb. of heads	numb. of experiments	\hat{p}
0	1	2	$\frac{1}{2}$
1	2	3	$\frac{2}{3}$
2	3	4	$\frac{3}{4}$
3	3	5	$\frac{3}{5}$
4	4	6	$\frac{2}{3}$
5	4	7	$\frac{4}{7}$

where time $t = 0$ represents prior data and their use is counted as the first two experiments (we inserted two measurements 1 and 0).

The evolution is




and it is much better then the previous case.

Finally, we can add not only two fictitious data but more, if we are convinced in our prior belief. Say that we decided to add five results 1 and five results 0 (for the probability $P = 0.5$). In our table, in the row 0, it will add 5 in the column “heads” and 10 in “results. So, we have

t	numb. of heads	numb. of experiments	\hat{p}
0	5	10	$\frac{1}{2}$
1	6	11	$\frac{6}{11}$
2	7	12	$\frac{7}{12}$
3	7	13	$\frac{7}{13}$
4	8	14	$\frac{4}{7}$
5	8	15	$\frac{8}{15}$

and the evolution is



and it respect relatively strongly our prior belief.

This example illustrates the commonly valid rules for inserting the prior belief into the on-line parameter estimation:

1. The ratio of the values of fictitious prior data express the values of the parameters.
2. The magnitude of the values determines the strength of our belief.

Program

```

#LyX 2.3 created this file. For more info see http://www.lyx.org/
\lyxformat 544
\begin_document
\begin_header
\save_transient_properties true
\origin unavailable
\textclass article
\begin_preamble
\usepackage{color}
\definecolor{red}{rgb}{0.7,0,0}
\end_preamble
\use_default_options true
\maintain_unincluded_children false
\language english
\language_package default
\inputencoding utf8
\fontencoding global
\font_roman "default" "default"
\font_sans "default" "default"
\font_typewriter "default" "default"
\font_math "auto" "auto"
\font_default_family default
\use_non_tex_fonts false
\font_sc false
\font_osf false
\font_sf_scale 100 100
\font_tt_scale 100 100
\use_microtype false
\use_dash_ligatures true
\graphics default
\default_output_format default
\output_sync 0
\bibtex_command default
\index_command default
\paperfontsize default
\spacing onehalf
\use_hyperref true
\pdf_bookmarks true
\pdf_bookmarksnumbered false
\pdf_bookmarksopen false

```

```

\pdf_bookmarksopenlevel 1
\pdf_breaklinks false
\pdf_pdfborder true
\pdf_colorlinks true
\pdf_backref false
\pdf_pdfusetitle true
\pdf_quoted_options "pdfstartview={XYZ 1000 1000 FitH}, pdfborder=0, colorlinks=true"
\papersize default
\use_geometry true
\use_package amsmath 1
\use_package amssymb 1
\use_package cancel 1
\use_package esint 1
\use_package mathdots 1
\use_package mathtools 1
\use_package mhchem 1
\use_package stackrel 1
\use_package stmaryrd 1
\use_package undertilde 1
\cite_engine basic
\cite_engine_type default
\biblio_style plain
\use_bibtopic false
\use_indices false
\paperorientation portrait
\suppress_date false
\justification true
\use_refstyle 1
\use_minted 0
\index Index
\shortcut idx
\color #008000
\end_index
\leftmargin 3cm
\topmargin 4cm
\rightmargin 4cm
\bottommargin 3cm
\secnumdepth 2
\tocdepth 2
\paragraph_separation skip

```

```

\defskip medskip
\is_math_indent 0
\math_numbering_side default
\quotes_style english
\dynamic_quotes 0
\papercolumns 1
\papersides 1
\paperpagestyle default
\tracking_changes false
\output_changes false
\html_math_output 0
\html_css_as_file 0
\html_be_strict false
\end_header

```

```

\begin_body

```

```

\begin_layout Subsubsection
Introduction
\end_layout

```

```

\begin_layout Standard
On-line (recursive) estimation consists in sequential update of the statistics
and if needed ongoing computation of th point estimates of the parameters.
Each step, the newly measured data are added and used for improving the
estimates.
The process starts with a prior statistics to be recomputed at the first
step.
If there is no prior information the prior statistics can be constructed
so they give uniform prior distribution of parameters (i.e.
each value of them i equally probable).
However, it is highly unreasonable, from the following reasons:
\end_layout

```

```

\begin_layout Enumerate
Always, there is some prior information - stronger or weaker, but it is.
\end_layout

```

```

\begin_layout Enumerate
With uniform prior density of parameters the initial estimates can be very

```

wild and for example if the model is used for control, they can be the cause of the failure of the whole control process.

\end_layout

\begin_layout Enumerate

At the beginning of estimation we use only few data (at the first step only one data record).

This record is under uncertainty and thus it can determine parameters that are far from the true values (imagine e.g. tossing a coin).

Then the estimates must be dragged back (with partially fixed statistics and they do not need to reach the true positions).

\end_layout

\begin_layout Standard

To prevent these inconveniences, we can

\end_layout

\begin_layout Itemize

set the initial statistics so that they produce reasonable point estimates of the parameters,

\end_layout

\begin_layout Itemize

tighten the initial estimates appropriately so that they hold on for a while and can be moved only by a larger portion of data.

\end_layout

\begin_layout Standard

These two steps are universal (for arbitrary model), but for each one the way how to do it is a bit different.

We shall demonstrate the way on the categorical model describing tossing of a coin.

\end_layout

\begin_layout Subsubsection

Example

\end_layout

```

\begin_layout Standard
Let us have a discrete variable
\begin_inset Formula  $y_t \in \left\{ 0,1 \right\}$  , $
\end_inset

\begin_inset Formula  $t=1,2,\cdots,N$ 
\end_inset

describing tossing a coin.
The value
\begin_inset Formula  $y_t=1$ 
\end_inset

represents the result
\begin_inset Quotes eld
\end_inset

head
\begin_inset Quotes erd
\end_inset

and
\begin_inset Formula  $y_t=0$ 
\end_inset

is
\begin_inset Quotes eld
\end_inset

tail
\begin_inset Quotes erd
\end_inset

.
We have tossed the coin 5 times ( $N=5$ ) and obtained the data
\end_layout

\begin_layout Standard
\align center

```


1 1 0 1 0
 \end{layout}

\begin{layout} Standard
 Perform step by step the estimation of the probability
 \begin{inset} Formula $P\left(\text{head}\right)$
 \end{inset}

.
 We know that in reality the the probability is
 \begin{inset} Formula $P=0.5$
 \end{inset}

.
 \end{layout}

\begin{layout} Standard
 The estimation can be performed according to the statistical definition
 of probability
 \begin{inset} Formula
 $\left[\right.$

$$P=\frac{\text{number of heads}}{\text{number of experiments}}$$

 $\left. \right]$
 \end{inset}

which is listed in the following table (
 \begin{inset} Formula \hat{p}
 \end{inset}

is the estimate of the probability)
 \end{layout}

\begin{layout} Standard
 \align center
 \begin{inset} Tabular
 \langle lyxtabular version="3" rows="6" columns="4">
 \langle features tabularvalignment="middle">
 \langle column alignment="center" valignment="top">
 \langle column alignment="center" valignment="top">

```

<column alignment="center" valignment="top">
<column alignment="center" valignment="top">
<row>
<cell alignment="center" valignment="top" bottomline="true" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
\begin_inset Formula  $t$ 
\end_inset

\end_layout

\end_inset
</cell>
<cell alignment="center" valignment="top" bottomline="true" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
numb.
  of heads
\end_layout

\end_inset
</cell>
<cell alignment="center" valignment="top" bottomline="true" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
numb.
  of experiments
\end_layout

\end_inset
</cell>
<cell alignment="center" valignment="top" bottomline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
\begin_inset Formula  $\hat{p}$ 

```

\end_inset

\end_layout

\end_inset

</cell>

</row>

<row>

<cell alignment="center" valignment="top" topline="true" rightline="true" usebox="none">

\begin_inset Text

\begin_layout Plain Layout

1

\end_layout

\end_inset

</cell>

<cell alignment="center" valignment="top" topline="true" rightline="true" usebox="none">

\begin_inset Text

\begin_layout Plain Layout

1

\end_layout

\end_inset

</cell>

<cell alignment="center" valignment="top" topline="true" usebox="none">

\begin_inset Text

\begin_layout Plain Layout

1

\end_layout

\end_inset

</cell>

<cell alignment="center" valignment="top" topline="true" leftline="true" rightline="true" usebox="non

\begin_inset Text

\begin_layout Plain Layout

```

1
\end_layout

\end_inset
</cell>
</row>
<row>
<cell alignment="center" valignment="top" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
2
\end_layout

\end_inset
</cell>
<cell alignment="center" valignment="top" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
2
\end_layout

\end_inset
</cell>
<cell alignment="center" valignment="top" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
2
\end_layout

\end_inset
</cell>
<cell alignment="center" valignment="top" leftline="true" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
1
\end_layout

```

```

\end_inset
</cell>
</row>
<row>
<cell alignment="center" valignment="top" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
3
\end_layout

\end_inset
</cell>
<cell alignment="center" valignment="top" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
2
\end_layout

\end_inset
</cell>
<cell alignment="center" valignment="top" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
3
\end_layout

\end_inset
</cell>
<cell alignment="center" valignment="top" leftline="true" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
\begin_inset Formula  $\frac{2}{3}$ 
\end_inset

```

```

\end_layout

\end_inset
</cell>
</row>
<row>
<cell alignment="center" valignment="top" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
4
\end_layout

\end_inset
</cell>
<cell alignment="center" valignment="top" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
3
\end_layout

\end_inset
</cell>
<cell alignment="center" valignment="top" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
4
\end_layout

\end_inset
</cell>
<cell alignment="center" valignment="top" leftline="true" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
\begin_inset Formula  $\frac{3}{4}$ 
\end_inset

```

\end_layout

\end_inset

</cell>

</row>

<row>

<cell alignment="center" valignment="top" rightline="true" usebox="none">

\begin_inset Text

\begin_layout Plain Layout

5

\end_layout

\end_inset

</cell>

<cell alignment="center" valignment="top" rightline="true" usebox="none">

\begin_inset Text

\begin_layout Plain Layout

3

\end_layout

\end_inset

</cell>

<cell alignment="center" valignment="top" usebox="none">

\begin_inset Text

\begin_layout Plain Layout

5

\end_layout

\end_inset

</cell>

<cell alignment="center" valignment="top" bottomline="true" leftline="true" rightline="true" usebox="

\begin_inset Text

\begin_layout Plain Layout

\begin_inset Formula $\frac{3}{5}$

\end_inset

```
\end_layout
```

```
\end_inset
```

```
</cell>
```

```
</row>
```

```
</lyxtabular>
```

```
\end_inset
```

```
\end_layout
```

```
\begin_layout Standard
```

We can see that at the beginning the estimates are totally wrong and only gradually they are refined.

The estimates evolution is

```
\end_layout
```

```
\begin_layout Standard
```

```
\align center
```

```
\begin_inset Graphics
```

```
filename est1.png
```

```
lyxscale 50
```

```
scale 20
```

```
\end_inset
```

```
\end_layout
```

```
\begin_layout Standard
```

Now, with our knowledge (that the coin is correct), we can construct so called fictitious prior data expressing our knowledge, e.g.

```
\end_layout
```

```
\begin_layout Standard
```

```
\align center
```



```

1 0
\end_layout

\begin_layout Standard
and use them before the estimation starts.
We obtain
\end_layout

\begin_layout Standard
\align center
\begin_inset Tabular
<lyxtabular version="3" rows="7" columns="4">
<features tabularvalignment="middle">
<column alignment="center" valignment="top">
<column alignment="center" valignment="top">
<column alignment="center" valignment="top">
<column alignment="center" valignment="top">
<row>
<cell alignment="center" valignment="top" bottomline="true" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
\begin_inset Formula  $t$ 
\end_inset

\end_layout

\end_inset
</cell>
<cell alignment="center" valignment="top" bottomline="true" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
numb.
of heads
\end_layout

\end_inset
</cell>

```

```

<cell alignment="center" valignment="top" bottomline="true" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
numb.
  of experiments
\end_layout

\end_inset
</cell>
<cell alignment="center" valignment="top" bottomline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
\begin_inset Formula  $\hat{p}$ 
\end_inset

\end_layout

\end_inset
</cell>
</row>
<row>
<cell alignment="center" valignment="top" topline="true" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
0
\end_layout

\end_inset
</cell>
<cell alignment="center" valignment="top" topline="true" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
1
\end_layout

```

```

\end_inset
</cell>
<cell alignment="center" valignment="top" topline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
2
\end_layout

\end_inset
</cell>
<cell alignment="center" valignment="top" topline="true" leftline="true" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
\begin_inset Formula  $\frac{1}{2}$ 
\end_inset

\end_layout

\end_inset
</cell>
</row>
<row>
<cell alignment="center" valignment="top" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
1
\end_layout

\end_inset
</cell>
<cell alignment="center" valignment="top" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
2
\end_layout

```

```

\end_inset
</cell>
<cell alignment="center" valignment="top" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
3
\end_layout

\end_inset
</cell>
<cell alignment="center" valignment="top" leftline="true" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
\begin_inset Formula  $\frac{2}{3}$ 
\end_inset

\end_layout

\end_inset
</cell>
</row>
<row>
<cell alignment="center" valignment="top" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
2
\end_layout

\end_inset
</cell>
<cell alignment="center" valignment="top" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
3

```

```

\end_layout

\end_inset
</cell>
<cell alignment="center" valignment="top" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
4
\end_layout

\end_inset
</cell>
<cell alignment="center" valignment="top" leftline="true" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
\begin_inset Formula  $\frac{3}{4}$ 
\end_inset

\end_layout

\end_inset
</cell>
</row>
<row>
<cell alignment="center" valignment="top" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
3
\end_layout

\end_inset
</cell>
<cell alignment="center" valignment="top" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout

```

```

3
\end_layout

\end_inset
</cell>
<cell alignment="center" valignment="top" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
5
\end_layout

\end_inset
</cell>
<cell alignment="center" valignment="top" leftline="true" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
\begin_inset Formula  $\frac{3}{5}$ 
\end_inset

\end_layout

\end_inset
</cell>
</row>
<row>
<cell alignment="center" valignment="top" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
4
\end_layout

\end_inset
</cell>
<cell alignment="center" valignment="top" rightline="true" usebox="none">
\begin_inset Text

```

```

\begin_layout Plain Layout
4
\end_layout

\end_inset
</cell>
<cell alignment="center" valignment="top" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
6
\end_layout

\end_inset
</cell>
<cell alignment="center" valignment="top" leftline="true" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
\begin_inset Formula  $\frac{2}{3}$ 
\end_inset

\end_layout

\end_inset
</cell>
</row>
<row>
<cell alignment="center" valignment="top" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
5
\end_layout

\end_inset
</cell>
<cell alignment="center" valignment="top" rightline="true" usebox="none">
\begin_inset Text

```

```
\begin_layout Plain Layout
```

```
4
```

```
\end_layout
```

```
\end_inset
```

```
</cell>
```

```
<cell alignment="center" valignment="top" usebox="none">
```

```
\begin_inset Text
```

```
\begin_layout Plain Layout
```

```
7
```

```
\end_layout
```

```
\end_inset
```

```
</cell>
```

```
<cell alignment="center" valignment="top" bottomline="true" leftline="true" rightline="true" usebox="
```

```
\begin_inset Text
```

```
\begin_layout Plain Layout
```

```
\begin_inset Formula  $\frac{4}{7}$ 
```

```
\end_inset
```

```
\end_layout
```

```
\end_inset
```

```
</cell>
```

```
</row>
```

```
</lyxtabular>
```

```
\end_inset
```

```
\end_layout
```

```
\begin_layout Standard
```

```
where time
```

```
\begin_inset Formula  $t=0$ 
```

```
\end_inset
```


represents prior data and their use is counted as the first two experiments
(we inserted two measurements 1 and 0).

\end_layout

\begin_layout Standard

The evolution is

\end_layout

\begin_layout Standard

\align center

\begin_inset Graphics

filename est2.png

lyxscale 50

scale 20

\end_inset

\end_layout

\begin_layout Standard

\align left

and it is much better then the previous case.

\end_layout

\begin_layout Standard

Finally, we can add not only two fictitious data but more, if we are convinced
in our prior belief.

Say that we decided to add five results 1 and five results 0 (for the probabili
ty

\begin_inset Formula $P=0.5$

\end_inset

).
In our table, in the row 0, it will add 5 in the column

\begin_inset Quotes eld

\end_inset

```

heads
\begin_inset Quotes erd
\end_inset

and 10 in
\begin_inset Quotes eld
\end_inset

results.
So, we have
\end_layout

\begin_layout Standard
\align center
\begin_inset Tabular
<lyxtabular version="3" rows="7" columns="4">
<features tabularvalignment="middle">
<column alignment="center" valignment="top">
<column alignment="center" valignment="top">
<column alignment="center" valignment="top">
<column alignment="center" valignment="top">
<row>
<cell alignment="center" valignment="top" bottomline="true" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
\begin_inset Formula  $t$ 
\end_inset

\end_layout

\end_inset
</cell>
<cell alignment="center" valignment="top" bottomline="true" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
numb.
of heads

```

```

\end_layout

\end_inset
</cell>
<cell alignment="center" valignment="top" bottomline="true" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
numb.
  of experiments
\end_layout

\end_inset
</cell>
<cell alignment="center" valignment="top" bottomline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
\begin_inset Formula  $\hat{p}$ 
\end_inset

\end_layout

\end_inset
</cell>
</row>
<row>
<cell alignment="center" valignment="top" topline="true" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
0
\end_layout

\end_inset
</cell>
<cell alignment="center" valignment="top" topline="true" rightline="true" usebox="none">
\begin_inset Text

```

```

\begin_layout Plain Layout
5
\end_layout

\end_inset
</cell>
<cell alignment="center" valignment="top" topline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
10
\end_layout

\end_inset
</cell>
<cell alignment="center" valignment="top" topline="true" leftline="true" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
\begin_inset Formula  $\frac{1}{2}$ 
\end_inset

\end_layout

\end_inset
</cell>
</row>
<row>
<cell alignment="center" valignment="top" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
1
\end_layout

\end_inset
</cell>
<cell alignment="center" valignment="top" rightline="true" usebox="none">
\begin_inset Text

```

```
\begin_layout Plain Layout
```

```
6
```

```
\end_layout
```

```
\end_inset
```

```
</cell>
```

```
<cell alignment="center" valignment="top" usebox="none">
```

```
\begin_inset Text
```

```
\begin_layout Plain Layout
```

```
11
```

```
\end_layout
```

```
\end_inset
```

```
</cell>
```

```
<cell alignment="center" valignment="top" leftline="true" rightline="true" usebox="none">
```

```
\begin_inset Text
```

```
\begin_layout Plain Layout
```

```
\begin_inset Formula  $\frac{6}{11}$ 
```

```
\end_inset
```

```
\end_layout
```

```
\end_inset
```

```
</cell>
```

```
</row>
```

```
<row>
```

```
<cell alignment="center" valignment="top" rightline="true" usebox="none">
```

```
\begin_inset Text
```

```
\begin_layout Plain Layout
```

```
2
```

```
\end_layout
```

```
\end_inset
```

```
</cell>
```

```
<cell alignment="center" valignment="top" rightline="true" usebox="none">
```

```

\begin_inset Text

\begin_layout Plain Layout
7
\end_layout

\end_inset
</cell>
<cell alignment="center" valignment="top" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
12
\end_layout

\end_inset
</cell>
<cell alignment="center" valignment="top" leftline="true" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
\begin_inset Formula  $\frac{7}{12}$ 
\end_inset

\end_layout

\end_inset
</cell>
</row>
<row>
<cell alignment="center" valignment="top" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
3
\end_layout

\end_inset
</cell>

```

```

<cell alignment="center" valignment="top" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
7
\end_layout

\end_inset
</cell>
<cell alignment="center" valignment="top" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
13
\end_layout

\end_inset
</cell>
<cell alignment="center" valignment="top" leftline="true" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
\begin_inset Formula  $\frac{7}{13}$ 
\end_inset

\end_layout

\end_inset
</cell>
</row>
<row>
<cell alignment="center" valignment="top" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
4
\end_layout

\end_inset

```

```

</cell>
<cell alignment="center" valignment="top" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
8
\end_layout

\end_inset
</cell>
<cell alignment="center" valignment="top" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
14
\end_layout

\end_inset
</cell>
<cell alignment="center" valignment="top" leftline="true" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
\begin_inset Formula  $\frac{4}{7}$ 
\end_inset

\end_layout

\end_inset
</cell>
</row>
<row>
<cell alignment="center" valignment="top" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
5
\end_layout

```



```

\end_inset
</cell>
<cell alignment="center" valignment="top" rightline="true" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
8
\end_layout

\end_inset
</cell>
<cell alignment="center" valignment="top" usebox="none">
\begin_inset Text

\begin_layout Plain Layout
15
\end_layout

\end_inset
</cell>
<cell alignment="center" valignment="top" bottomline="true" leftline="true" rightline="true" usebox="
\begin_inset Text

\begin_layout Plain Layout
\begin_inset Formula  $\frac{8}{15}$ 
\end_inset

\end_layout

\end_inset
</cell>
</row>
</lyxtabular>

\end_inset

\end_layout

```

```

\begin_layout Standard
and the evolution is
\end_layout

```

```

\begin_layout Standard
\align center
\begin_inset Graphics
filename est3.png
lyxscale 50
scale 20

```

```

\end_inset

```

```

\end_layout

```

```

\begin_layout Standard
and it respect relatively strongly our prior belief.
\end_layout

```

```

\begin_layout Standard
This example illustrates the commonly valid rules for inserting the prior
belief into the on-line parameter estimation:
\end_layout

```

```

\begin_layout Enumerate
The ratio of the values of fictitious prior data express the values of the
parameters.
\end_layout

```

```

\begin_layout Enumerate
The magnitude of the values determines the strength of our belief.
\end_layout

```

```

\begin_layout Standard
Program
\end_layout

```

```

\begin_layout Standard

```

```

\begin_inset CommandInset include
LatexCommand verbatiminput
filename "d2_init0.lyx"

```

```

\end_inset

```

```

\end_layout

```

```

\begin_layout Subsubsection
\begin_inset Note Note
status collapsed

```

```

\begin_layout Subsubsection
Inserting prior belief for estimation of expectation
\end_layout

```

```

\begin_layout Plain Layout
For estimation of the expectation (i.e.
  for models exponential, Bernoulli, binomial and similar ones), the statistics
  (sum and counter) and their update are
\begin_inset Formula
\[

$$S_{\{t\}}=S_{\{t-1\}}+y_{\{t\}}$$

\]

```

```

\end_inset

```

```

\begin_inset Formula
\[

$$\kappa_{\{t\}}=\kappa_{\{t-1\}}+1$$

\]

```

```

\end_inset

```

```

\end_layout

```

```

\begin_layout Plain Layout

```

Here we can proceed as follows: let

```
\begin_inset Formula $p_{0}$
\end_inset
```

is our guess of the parameter.

Then set

```
\begin_inset Formula
\[
\kappa_{0}=N_{0}
\]
```

```
\end_inset
```

```
\begin_inset Formula
\[
S_{0}=p_{0}\kappa_{0}
\]
```

```
\end_inset
```

where

```
\begin_inset Formula $N_{0}$
\end_inset
```

is the number of fictitious data expressing the strength of our belief
in the value

```
\begin_inset Formula $p_{0}$
\end_inset
```

.

```
\end_layout
```

```
\begin_layout Plain Layout
```

Then

```
\begin_inset Formula
```

```
\[
\hat{p}=\frac{S_{0}}{\kappa_{0}}=\frac{p_{0}\kappa_{0}}{\kappa_{0}}=p_{0}
\]
```

`\end_inset`

independently of

`\begin_inset Formula N_{0}`

`\end_inset`

which is free for setting the strength.

`\end_layout`

`\begin_layout Subsubsection`

Inserting prior belief for regression models

`\end_layout`

`\begin_layout Plain Layout`

For scalar regression model it holds

`\begin_inset Formula`

`\[`

$$V_t = V_{t-1} + \Psi_t \Psi_t^{\prime}$$

`\]`

`\end_inset`

where

`\begin_inset Formula $\Psi_t = \left[y_t, \psi_t^{\prime} \right]^{\prime}$`

`\end_inset`

is extended regression vector with

`\begin_inset Formula ψ_t`

`\end_inset`

being the regression vector.

The point estimate is

`\begin_inset Formula`

`\[`

$$\hat{\theta} = V_{\psi}^{-1} V_{y\psi}$$

`\]`

`\end_inset`

with the partitioning

```

\begin_inset Formula
\[
V=\left[\begin{array}{cc}
V_{\{y\}} & V_{\{y\psi}^{\{'}}\\
V_{\{y\psi\}} & V_{\{\psi\}}
\end{array}\right]
\end_inset

```

```

\end_inset

```

with the dimensions given by the extended regression vector.

```

\end_layout

```

```

\begin_layout Plain Layout
So, if we want the parameters to be
\begin_inset Formula $\theta_0$
\end_inset

```

then we set

```

\end_layout

```

```

\begin_layout Plain Layout
\begin_inset Formula
\[
\tilde{V}=\left[\begin{array}{cc}
1 & \theta_0^{\{'}}\\
\theta_0 & I
\end{array}\right]
\end_inset

```

```

\end_inset

```

with

```

\begin_inset Formula $I$
\end_inset

```

being a unit matrix, and the initial statistics are

```

\begin_inset Formula
\[
\kappa_0=N_0

```

\]

\end_inset

\begin_inset Formula

\[

$V_0 = \kappa_0 \tilde{V}$

\]

\end_inset

\end_layout

\begin_layout Plain Layout

Then we obtain

\begin_inset Formula

\[

$\hat{\theta} = V_{\psi}^{-1} V_{y\psi} = \left(\kappa_0 I \right)^{-1} \times \kappa_0 \theta = \theta$

\]

\end_inset

and again

\begin_inset Formula N_0

\end_inset

express our belief in the estimate

\begin_inset Formula θ

\end_inset

\end_layout

\begin_layout Plain Layout

Program

\end_layout

\begin_layout Plain Layout

```
\begin_inset CommandInset include
LatexCommand verbatiminput
filename "Test4b.sce"

\end_inset

\end_layout

\end_inset

\end_layout

\end_body
\end_document
```